



Moutard type transform for matrix generalized analytic functions and gauge transforms

Roman Novikov, Iskander Taimanov

► To cite this version:

Roman Novikov, Iskander Taimanov. Moutard type transform for matrix generalized analytic functions and gauge transforms. Russian Mathematical Surveys, 2016, 71 (5), pp.970-972. hal-01348015

HAL Id: hal-01348015

<https://hal.science/hal-01348015>

Submitted on 22 Jul 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Moutard type transform for matrix generalized analytic functions and gauge transforms *

R.G. Novikov [†] I.A. Taimanov [‡]

Considerable progress in the theory of Darboux-Moutard type transforms for two-dimensional linear differential systems with applications to geometry, spectral theory, and soliton equations has been achieved recently, see, e.g., [1, 2, 3, 4]. In the present note we derive such a transformation for the matrix generalized function system

$$\partial_{\bar{z}}\Psi + A\Psi + B\bar{\Psi} = 0, \quad (1)$$

where $\partial_{\bar{z}} = \frac{\partial}{\partial \bar{z}}$, the coefficients A and B and solutions Ψ are $(N \times N)$ -matrix functions on D , with D an open simply connected domain in \mathbb{C} . In particular, this generalizes the transform for $N = 1$ found in [4] with $A = 0$. In addition, we show that the Moutard type transform for system (1) with $B = 0$ is equivalent to a gauge transform for the connection $\nabla_{\bar{z}} = \partial_{\bar{z}} + A$. In turn, our studies show that the Moutard type transform for system (1) with $A = 0$ can be treated as a proper analog of the aforementioned gauge transform.

As for $N = 1$, system (1) is reduced to the system

$$\partial_{\bar{z}}\Psi + B\bar{\Psi} = 0, \quad (2)$$

i.e. to system (1) with $A = 0$, by the gauge transform

$$\Psi \rightarrow \tilde{\Psi} = g^{-1}\Psi, \quad B \rightarrow \tilde{B} = g^{-1}Bg, \quad \partial_{\bar{z}}g + Ag = 0, \quad \det g \neq 0.$$

We say that the system

$$\partial_z\Psi^+ - \bar{\Psi}^+B = 0 \quad (3)$$

is conjugate to system (2) (see [5] for a similar definition for $N = 1$).

We have the following result.

*The work was done during the visit of the second author (I.A.T.) to Centre de Mathématiques Appliquées of École Polytechnique in July 2016 and was supported by RSF (grant 14-11-00441).

[†]CNRS (UMR 7641), Centre de Mathématiques Appliquées, École Polytechnique, 91128 Palaiseau, France, and IEPT RAS, 117997 Moscow, Russia; e-mail: novikov@cmap.polytechnique.fr

[‡]Sobolev Institute of Mathematics, 630090 Novosibirsk, Russia, and Novosibirsk State University, 630090 Novosibirsk, Russia; e-mail: taimanov@math.nsc.ru

Theorem 1 *Systems (2) and (3) are covariant, i.e. mapped into the systems of the same type, with respect to the Moutard type transform*

$$\begin{aligned}\Psi &\rightarrow \tilde{\Psi} = \Psi - F \omega_{F,F^+}^{-1} \omega_{\Psi,F^+}, \\ \Psi^+ &\rightarrow \tilde{\Psi}^+ = \Psi^+ - \omega_{F,\Psi^+} \omega_{F,F^+}^{-1} F^+, \\ B &\rightarrow \tilde{B} = B + F \omega_{F,F^+}^{-1} F^+, \end{aligned} \quad (4)$$

where F and F^+ are arbitrary fixed solutions of (2) and (3), respectively,

$$\partial_{\bar{z}} \omega_{\Phi,\Phi^+} = \Phi^+ \bar{\Phi}, \quad \text{Re } \omega_{\Phi,\Phi^+} = 0, \quad (5)$$

for Φ and Φ^+ meeting equations (2) and (3), and $\det \omega_{F,F^+} \neq 0$.

For finding ω_{Φ,Φ^+} satisfying (5) we use also that $\partial_z \omega_{\Phi,\Phi^+} = -\bar{\Phi}^+ \Phi$. In addition, our definition of ω_{Φ,Φ^+} is self-consistent up to a pure imaginary matrix integration constant in view of the identity $\partial_z \Phi^+ \bar{\Phi} = -\partial_{\bar{z}} \bar{\Phi}^+ \Phi$. The latter equality follows from systems (2) and (3) for Φ and Φ^+ , respectively. We recall that the domain D is simply connected.

Given ω_{F,F^+} , ω_{Ψ,F^+} , and ω_{F,Ψ^+} , Theorem 1 is proved by straightforward computations.

In addition, for the system

$$\partial_{\bar{z}} \Psi + A \Psi = 0, \quad (6)$$

i.e., for system (1) with $B = 0$, the following result also holds.

Proposition 1 *System (6) is covariant under the following Moutard type transform*

$$\Psi \rightarrow \tilde{\Psi} = \Psi - F \hat{\omega}_{F,F^+}^{-1} \hat{\omega}_{\Psi,F^+}, \quad A \rightarrow \tilde{A} = A + F \hat{\omega}_{F,F^+}^{-1} F^+, \quad (7)$$

where F is an arbitrary fixed solution of (6), F^+ is an arbitrary fixed matrix function,

$$\partial_{\bar{z}} \hat{\omega}_{\Phi,F^+} = F^+ \Phi \quad (8)$$

for any matrix function Φ , and $\det \hat{\omega}_{F,F^+} \neq 0$.

Equations (7) and (8) are analogs of equations (4) and (5). However, in difference with (5), we do not require that the matrix functions $\hat{\omega}_{F,F^+}$ would be pure imaginary. Equation (8) is solvable for $\hat{\omega}_{F,F^+}$ and Proposition 1 is proved by straightforward computations.

REMARK. Let $A, \tilde{A}, \Psi, F, F^+$, and $\hat{\omega}_{\Phi,F^+}$ be the same as in Proposition 1. Let

$$g = 1 - F \hat{\omega}_{F,F^+}^{-1} \Lambda, \quad \Lambda_{\bar{z}} = \Lambda A + F^+.$$

Then

$$\partial_{\bar{z}}(g\Psi) + \tilde{A}(g\Psi) = 0.$$

It is proved by straightforward computations and it shows that for invertible g the transform $A \rightarrow \tilde{A}$ reduces to a gauge transform.

References

- [1] Delong Yu, Q.P. Liu, and Shikun Wang: Darboux transformation for the modified Veselov-Novikov equation. *J. Phys. A* **35**:16 (2002), 3779–3785.
- [2] I.A. Taimanov and S.P.Tsarev: Two-dimensional rational solitons and their blowup via the Moutard transformation. *Theoret. and Math. Phys.* **157**:2 (2008), 1525–1541.
- [3] I.A. Taimanov: The Moutard transformation of two-dimensional Dirac operators and Möbius geometry. *Math. Notes* **97** (2015), 124–135.
- [4] P.G. Grinevich and R.G. Novikov: Generalized analytic functions, Moutard-type transforms, and holomorphic maps. *Functional Analysis and its Appl.* **50**:2 (2016), 150–152.
- [5] I.N. Vekua: *Generalized Analytic Functions*. Pergamon Press, London–Paris–Frankfurt, 1962.